arXiv:1603.08184v1 [math.GR] 27 Mar 2016

Permutation-like Matrix Groups with a Maximal Cycle of Length Power of Two

Guodong Deng, Yun Fan

School of Mathematics and Statistics
Central China Normal University, Wuhan 430079, China

Abstract

If every element of a matrix group is similar to a permutation matrix, then it is called a permutation-like matrix group. References [4], [5] and [6] showed that, if a permutation-like matrix group contains a maximal cycle such that the maximal cycle generates a normal subgroup and the length of the maximal cycle equals to a prime, or a square of a prime, or a power of an odd prime, then the permutation-like matrix group is similar to a permutation matrix group. In this paper, we prove that if a permutation-like matrix group contains a maximal cycle such that the maximal cycle generates a normal subgroup and the length of the maximal cycle equals to any power of 2, then it is similar to a permutation matrix group.

Key words: permutation-like matrix group, permutation matrix group, maximal cycle.

MSC2010: 15A18, 15A30, 20H20.

1 Introduction

By $\mathrm{GL}_d(\mathbb{C})$ we denote the complex general linear group of dimension (degree) d consisting of invertible $d \times d$ complex matrices. Any subgroup $\mathcal{G}$ of $\mathrm{GL}_d(\mathbb{C})$ is said to be a *matrix group* of dimension d. If there is a $T \in \mathrm{GL}_d(\mathbb{C})$ such that for every $A \in \mathcal{G}$ the conjugation $T^{-1}AT$ is a permutation matrix, then we say that $\mathcal{G}$ is a *permutation matrix group*. If for any $A \in \mathcal{G}$ there is a $T \in \mathrm{GL}_d(\mathbb{C})$ such that $T^{-1}AT$ is a permutation matrix, then $\mathcal{G}$ is called a *permutation-like matrix group*.

Cigler [3, 4] showed that a permutation-like matrix group is not a permutation matrix group in general. A $d \times d$ matrix is called a *maximal cycle* if it

Email address: yfan@mail.ccnu.edu.cn (Yun Fan).

is similar to a permutation matrix corresponding to the cycle permutation of length d. Cigler conjectured that:

Conjecture: *If a permutation-like matrix group contains a maximal cycle, then it is a permutation matrix group.*

Let $\mathcal{G}$ be a permutation-like matrix group containing a maximal cycle which generates a normal cyclic subgroup. Cigler [3, 4] proved that if the dimension of $\mathcal{G}$ is a prime then $\mathcal{G}$ is a permutation matrix group. In [5, 6], we further proved that the $\mathcal{G}$ is a permutation matrix group if the dimension is a square of a prime, or any power of an odd prime.

In this paper we prove:

Theorem 1.1. *Let $\mathcal{G}$ be a permutation-like matrix group of dimension 2^n where n is any positive integer. If $\mathcal{G}$ contains a maximal cycle which generates a normal cyclic subgroup, then $\mathcal{G}$ is a permutation matrix group.*

Necessary preparations for the proof of the theorem are made in Section 2. For fundamentals of the group theory, please refer to [1, 7]. In Section 3 we prove the theorem in the case when the quotient group of $\mathcal{G}$ over the cyclic subgroup generated by the maximal cycle is a cyclic group, see Theorem 3.9 below. In Section 4 we prove the theorem for the case when the quotient group is not cyclic, see Theorem 4.3 below.

2 Preparations

We sketch necessary fundamentals from group theory, number theory and matrix theory, and formulate them in a way suitable to our later quotations.

The order of an element C of a group $\mathcal{G}$ is denoted by $\mathrm{ord}_{\mathcal{G}}(C)$, or $\mathrm{ord}(C)$ in short if the group is known from context. By $\langle C\rangle$ we denote the cyclic group generated by C. Assume that $\mathrm{ord}(C) = d$. Then $\langle C\rangle$ is isomorphic to the additive group of the residue ring $\mathbb{Z}_d$ of the integer ring $\mathbb{Z}$ modulo d by mapping $j \in \mathbb{Z}_d$ to $C^j \in \langle C\rangle$. In this way, the properties of the cyclic group $\langle C\rangle$ are exactly corresponding to the properties of the additive group $\mathbb{Z}_d$, some of which are listed as follows.

- $\langle C^r\rangle = \langle C\rangle$ if and only if $r \in \mathbb{Z}_d^*$, where $\mathbb{Z}_d^*$ denotes the multiplicative group consisting of the reduced residue classes in $\mathbb{Z}_d$; in that case C^r is said to be a *generator* of the cyclic group $\langle C\rangle$.
- Each automorphism α of $\langle C\rangle$ is corresponding to exactly one $r \in \mathbb{Z}_d^*$ such that $\alpha(C^j) = C^{rj}$ for any $C^j \in \langle C\rangle$, and μ_r is an automorphism of the additive group $\mathbb{Z}_d$, where $\mu_r(j) = rj$ for any $j \in \mathbb{Z}_d$.
- If $\langle C\rangle$ is normal in $\mathcal{G}$, then $\mathcal{G}/\mathrm{Cent}_{\mathcal{G}}(\langle C\rangle)$ is homomorphic to a subgroup of $\mathbb{Z}_d^*$, where $\mathrm{Cent}_{\mathcal{G}}(\langle C\rangle) = \{A \in \mathcal{G} \mid A^{-1}CA = C\}$, called the *centralizer* of $\langle C\rangle$ in $\mathcal{G}$. If $\mathrm{Cent}_{\mathcal{G}}(\langle C\rangle) \subseteq \langle C\rangle$, then $\langle C\rangle$ is said to be *self-centralized* in $\mathcal{G}$.

In this paper we are concerned with the cyclic group of order 2^n (with $\mathbb{Z}_{2^n}$ correspondingly). The next lemma has appeared in [6].

Lemma 2.1. *Assume that $\mathcal{G} = \langle C \rangle$ is a cyclic group of order $|\mathcal{G}| = 2^n$ and $0 \le a < n$. Let $\mathcal{G}^{2^a} = \{C^{2^a t} \mid 0 \le t < 2^{n-a}\}$. Then:*

(i) *$\mathcal{G}^{2^a} = \langle C^{2^a} \rangle$ is a cyclic subgroup of $\mathcal{G}$ of order 2^{n-a} generated by C^{2^a}.*

(ii) *The mapping $C^j \mapsto C^{2^a j}$ is a homomorphism of $\mathcal{G}$ onto $\mathcal{G}^{2^a}$ with kernel $\mathcal{G}^{2^{n-a}}$. Hence, for any generator G of the cyclic group $\mathcal{G}^{2^a}$ there are exactly 2^a generators of $\mathcal{G}$ which are mapped to G.* □

By $\Phi_m(x)$ we denote the cyclotomic polynomial of degree m, i.e., $\Phi_m(x) = \prod_\theta (x - \theta)$ with θ running over the primitive m-th roots of unity. Note that Lemma 2.1 (ii) (with $k = n - a$) implies that

$$\prod_{\omega} (x^{2^a} - \omega) = \Phi_{2^{k+a}}(x) \tag{2.1}$$

where the subscript ω of $\prod$ runs over the primitive 2^k-th roots of unity. In the specific case when $k = 1$, since -1 is the unique primitive 2-th root of unity, we get a known formula: $\Phi_{2^{a+1}}(x) = x^{2^a} + 1$.

Remark 2.2. The structure of the multiplicative group $\mathbb{Z}_{2^n}^*$ is known (e.g., see [2, Remark 9]). If $n \ge 3$ then the following hold:

(i) $\mathbb{Z}_{2^n}^* = \langle 5 \rangle \times \langle -1 \rangle$ with $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(5) = 2^{n-2}$ and $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(-1) = 2$.

(ii) $\langle 5 \rangle = \{1\} \cup \{1 + 2^b v \mid b = 2, 3, \cdots, n-1,\ v = 1, 3, \cdots, 2^{n-b} - 1\}$, and $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(1 + 2^b v) = 2^{n-b}$.

(iii) $-\langle 5 \rangle = \{-1\} \cup \{-1 + 2^b v \mid b = 2, 3, \cdots, n-1,\ v = 1, 3, \cdots, 2^{n-b} - 1\}$, and $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(-1 + 2^b v) = 2^{n-b}$.

In our study, the behaviors of the elements -1, $1 + 2^b v$ and $-1 + 2^b v$ are quite different. But note that $(-1 + 2^b v)^\ell \in \langle 5 \rangle$ provided ℓ is even.

For the subgroups of $\mathbb{Z}_{2^n}^*$, we have the following result.

Lemma 2.3. *If $\mathcal{H}$ is a subgroup of the multiplicative group $\mathbb{Z}_{2^n}^*$ where $n \ge 3$, then one of the following holds:*

(i) *$\mathcal{H}$ is a cyclic group, and, if the order $|\mathcal{H}| = 2^a$, then $0 \le a \le n - 2$ and $\mathcal{H} = \langle 1 + 2^{n-a} \rangle$, or $\mathcal{H} = \langle -1 + 2^{n-a} \rangle$, or $\mathcal{H} = \langle -1 \rangle$ (hence $a = 1$).*

(ii) *$\mathcal{H} = \langle -1 \rangle \times \langle 1 + 2^{n-a} \rangle$ with $1 \le a \le n - 2$.*

Proof. By Remark 2.2, we have a surjective homomorphism $\rho : \mathbb{Z}_{2^n}^* \to \langle 5\rangle$ such that $\rho(-1) = 1$, $\rho(1 + 2^b v) = 1 + 2^b v$ and $\rho(-1 + 2^b v) = 1 + 2^b(2^{n-b} - v)$. Then $\rho(\mathcal{H})$ is a subgroup of $\langle 5\rangle$, hence $\rho(\mathcal{H})$ is a cyclic group of order 2^a with $0 \le a \le n-2$. If $a = 0$, then $\mathcal{H} = 1$ or $\mathcal{H} = \langle -1\rangle$. Assume that $a \ge 1$ in the following. Take a $k \in \mathcal{H}$ such that $\rho(k)$ is a generator of $\rho(\mathcal{H})$. Then $\mathcal{H} = \langle k\rangle \times (\mathcal{H} \cap \langle -1\rangle)$, and $k = 1+2^{n-a}$ or $k = -1+2^{n-a}$. If $\mathcal{H} \cap \langle -1\rangle = 1$, then (i) holds. Otherwise, $\mathcal{H} \cap \langle -1\rangle = \langle -1\rangle$, hence $\mathcal{H} = \langle k\rangle \times \langle -1\rangle$; if $k = -1+2^{n-a}$, we can replace k by $-k$. In a word, (ii) holds. □

For any non-zero integer k, by $\nu_2(k)$ we denote the 2-adic valuation of k, i.e., $k = 2^{\nu_2(k)}u$ for an odd integer u.

Lemma 2.4. *Let $r \in \mathbb{Z}_{2^n}^*$ such that $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$ with $1 \le a \le n-2$.*

(i) *If $r = -1$, then $a = 1$ and $r^{2^a-1} + r^{2^a-2} + \cdots + r + 1 = 0$.*

(ii) *If $r = 1 + 2^{n-a}v$ with v being odd, then*

$$\nu_2(r^{2^a-1} + r^{2^a-2} + \cdots + r + 1) = a.$$

(iii) If $r = -1 + 2^{n-a}v$ with v being odd, then

$$\nu_2(r^{2^a-1} + r^{2^a-2} + \cdots + r + 1) = n - 1.$$

Proof. (i) is obvious.

(ii). Since $\nu_2(2^a - j) = \nu_2(j)$ for $0 < j < 2^a$, for the binomial coefficients $\binom{2^a}{k}$ we get that

$$\nu_2\binom{2^a}{k} = \nu_2\Big(2^a \cdot \frac{2^a-1}{1} \cdot \cdots \cdot \frac{2^a-(k-1)}{k-1} \cdot \frac{1}{k}\Big) = a - \nu_2(k). \tag{2.2}$$

So

$$r^{2^a-1} + \cdots + r + 1 = \frac{r^{2^a}-1}{r-1} = \frac{(1+2^{n-a}v)^{2^a}-1}{2^{n-a}v} \equiv 2^a \pmod{2^{a+1}}.$$

(iii). Noting that $r - 1 = 2u$ with u being odd, by Eqn (2.2) we obtain that

$$r^{2^a-1} + \cdots + r + 1 = \frac{r^{2^a}-1}{r-1} = \frac{(-1+2^{n-a}v)^{2^a}-1}{2u} \equiv 2^{n-1}v' \pmod{2^n};$$

where $v' \equiv -vu^{-1} \pmod{2^n}$. □

Lemma 2.5. *Let $\mathcal{G} = \langle A, C\rangle$ be a finite group generated by A and C. If $\langle C\rangle$ is a self-centralized normal cyclic subgroup of order 2^n and $A^{-1}CA = C^r$ for an $r \in \mathbb{Z}_{2^n}$ with $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$, then $|\mathcal{G}| = 2^{n+a}$ and:*

(i) *If $r \not\equiv -1 \pmod{2^n}$, then there is an $A' \in \mathcal{G}$ such that $\mathcal{G} = \langle A', C\rangle$ with $A'^{-1}CA' = C^r$ and $A'^{2^a} = 1$.*

(ii) *If* $r \equiv -1 \pmod{2^n}$ *(hence* $a = 1$*), then one of the following two holds:*

(ii.1) $A^2 = 1$*; in that case* $(AC^k)^2 = 1$ *for any* $C^k \in \langle C\rangle$ *(i.e.,* $\mathcal{G}$ *is the dihedral group of order* 2^{n+1}*).*

(ii.2) $A^2 = C^{2^{n-1}}$*; in that case* $(AC^k)^2 = C^{2^{n-1}}$ *for any* $C^k \in \langle C\rangle$ *(i.e.,* $\mathcal{G}$ *is the generalized quaternion group of order* 2^{n+1}*).*

Proof. Since $\langle C\rangle$ is self-centralized, the quotient group $\mathcal{G}/\langle C\rangle \cong \langle A\rangle/\langle A\rangle\cap\langle C\rangle$ is isomorphic to a cyclic subgroup $\langle r\rangle$ of the group $\mathbb{Z}_{2^n}^*$. So $|\langle A\rangle/\langle A\rangle \cap \langle C\rangle| = 2^a$, $|\mathcal{G}| = 2^{n+a}$ and $A^{2^a} \in \langle C\rangle$. Let $\langle C\rangle^A = \{C^t \mid A^{-1}C^tA = C^t\}$, i.e., the centralizer in $\langle C\rangle$ of A. Then $A^{2^a} \in \langle C\rangle^A$. Since $A^{-1}C^tA = C^{tr}$,

$$A^{-1}C^tA = C^t \quad \text{iff} \quad C^{tr-t} = 1 \quad \text{iff} \quad t(r-1) \equiv 0 \pmod{2^n}. \tag{2.3}$$

Suppose that $r = -1 + 2^{n-a}v$ where v is zero (i.e., $r = -1$) or odd. Since $n - a \geq 2$, we have $r - 1 = -2 + v2^{n-a} = 2u$ with u being odd.

$$A^{-1}C^tA = C^t \quad \text{iff} \quad 2ut \equiv 0 \pmod{2^n} \quad \text{iff} \quad t \equiv 0 \pmod{2^{n-1}}.$$

Thus $\langle C\rangle^A = \langle C^{2^{n-1}}\rangle$ which is a group of order 2 consisting of two elements $\{1, C^{2^{n-1}}\}$. So

$$A^{2^a} \in \langle C^{2^{n-1}}\rangle = \{1, C^{2^{n-1}}\}.$$

In particular, if $v = 0$, i.e., $r = -1$, then we obtain the conclusion (ii).

Next, we assume that $r = -1 + 2^{n-a}v$ with v being odd. Then $A^{2^a} = 1$ or $A^{2^a} = C^{2^{n-1}}$. If $A^{2^a} = 1$ then we are done. Suppose that $A^{2^a} = C^{2^{n-1}}$. Note that for any integer k and any positive integer j, by a direct computation we have the following formula:

$$(AC^k)^j = (AC^k)\cdots(AC^k)(AC^k) = A^jC^{k(r^{j-1}+\cdots+r+1)}. \tag{2.4}$$

By Lemma 2.4(iii), $r^{2^a-1} + \cdots + r + 1 = 2^{n-1}u$ with u being odd. We have an integer u' such that $u'u \equiv 1 \pmod{2^n}$. Taking $j = 2^a$ and $k = u'$ in Eqn (2.4), we obtain

$$(AC^{u'})^{2^a} = A^{2^a}C^{u'(r^{2^a-1}+\cdots+r+1)} = A^{2^a}C^{2^{n-1}u'u} = C^{2^{n-1}}C^{2^{n-1}} = 1.$$

Replacing A by $A' = AC^{u'}$, we get that $\mathcal{G} = \langle A', C\rangle$, $A'^{-1}CA' = C^r$ and $A'^{2^a} = 1$. So (i) holds.

Finally, assume that $r = 1 + 2^{n-a}v$ with v being odd. Then

$$t(r-1) \equiv 0 \pmod{2^n} \quad \text{iff} \quad 2^{n-a}vt \equiv 0 \pmod{2^n} \quad \text{iff} \quad t \equiv 0 \pmod{2^a}.$$

By Eqn (2.3), $\langle C\rangle^A = \langle C^{2^a}\rangle$; hence $A^{2^a} \in \langle C\rangle^A = \langle C^{2^a}\rangle$. So, we can find an integer h such that $A^{2^a}C^{2^ah} = 1$. By Lemma 2.4(ii) we can assume that $r^{2^a-1} + \cdots + r + 1 = 2^au$ with u being odd. We have an integer u' such that $u'u \equiv 1 \pmod{2^n}$, Taking $j = 2^a$ and $k = u'h$ in Eqn (2.4), we obtain

$$(AC^{u'h})^{2^a} = A^{2^a}C^{u'h(r^{2^a-1}+\cdots+r+1)} = A^{2^a}C^{2^auu'h} = 1.$$

Replacing A by $A' = AC^{u'h}$, we get the conclusion (i). □

The next lemma shows that the condition "self-centralized" is usually satisfied in our study.

Lemma 2.6 ([4, Proposition 4.2]). *If $\mathcal{G}$ is a permutation-like matrix group and $C \in \mathcal{G}$ is a maximal cycle, then $\langle C \rangle$ is self-centralized in $\mathcal{G}$.*

There is also a proof of this lemma in [6, Lemma 2.7].

For any complex matrix A, by $\mathrm{char}_A(x)$ we denote the characteristic polynomial of A, i.e., $\mathrm{char}_A(x) = \det(xI - A)$, where I denotes the identity matrix.

Lemma 2.7 ([5, Lemma 2.1], [6, Lemma 2.3]). *The following two are equivalent to each other:*

(i) *A is similar to a permutation matrix.*

(ii) *A is diagonalizable and $\mathrm{char}_A(x) = \prod_i (x^{\ell_i} - 1)$.*

If it is the case, then each factor $x^{\ell_i} - 1$ of $\mathrm{char}_A(x)$ corresponds to exactly one ℓ_i-cycle of the cycle decomposition of the permutation of the permutation matrix. □

Note that $x^n - 1 = \prod_{k|n} \Phi_k(x)$, where $\Phi_k(x)$ is the cyclotomic polynomial of degree k, and "$k|n$" stands for that k divides n. We get an immediate consequence of the above lemma as follows.

Corollary 2.8. *Let A be a matrix similar to a permutation matrix, and m, n be positive integers. If $\Phi_n(x)^m \,\big|\, \mathrm{char}_A(x)$, then $\Phi_k(x)^m \,\big|\, \mathrm{char}_A(x)$ for any $k \,|\, n$.* □

3 The quotient $\mathcal{G}/\langle C \rangle$ is cyclic

In this section we always assume:

- $A, C \in \mathrm{GL}_{2^n}(\mathbb{C})$;
- C is a maximal cycle;
- $A^{-1}CA = C^r$, where $r \in \mathbb{Z}_{2^n}^*$ and $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$ with $0 \le a \le n-2$.

Let λ be a primitive 2^n-th root of unity. Then

$$\{\lambda^j \mid j \in \mathbb{Z}_{2^n}\} = \{1 = \lambda^0, \lambda, \lambda^2, \cdots, \lambda^{2^n-1}\} \tag{3.1}$$

is the spectrum (i.e., the set of eigenvalues) of C. The eigen-subspace of each eigenvalue λ^j of C, denoted by $\mathrm{E}(\lambda^j)$, has dimension 1; and the complex space $\mathbb{C}^{2^n}$ is decomposed into a direct sum of the eigen-subspaces:

$$\mathbb{C}^{2^n} = \bigoplus_{j \in \mathbb{Z}_{2^n}} \mathrm{E}(\lambda^j) = \mathrm{E}(\lambda^0) \oplus \mathrm{E}(\lambda) \oplus \mathrm{E}(\lambda^2) \oplus \cdots \oplus \mathrm{E}(\lambda^{2^n-1}). \tag{3.2}$$

Taking any non-zero vector $e_j \in \mathrm{E}(\lambda^j)$ for every index $j \in \mathbb{Z}_{2^n}$, we get a basis of $\mathbb{C}^{2^n}$:

$$\mathcal{E} = \{e_j \mid j \in \mathbb{Z}_{2^n}\} = \{e_0, e_1, e_2, \cdots, e_{2^n-1}\}, \tag{3.3}$$

which will play a key role in our study. For convenience we denote the diagonal blocked matrix $\begin{pmatrix} B_1 & & \\ & \ddots & \\ & & B_m \end{pmatrix}$ by $B_1 \oplus \cdots \oplus B_m$. With the basis $\mathcal{E}$ in Eqn (3.3), the transformation C on $\mathbb{C}^{2^n}$ has the diagonal matrix

$$C|_{\mathcal{E}} = 1 \oplus \lambda \oplus \lambda^2 \oplus \cdots \oplus \lambda^{2^n-1}.$$

For the above, please see [6, Lemma 2.4].

Consider the matrix $A|_{\mathcal{E}}$ with respect to the basis $\mathcal{E}$ in Eqn (3.3). Recall that $A^{-1}CA = C^r$ where $r \in \mathbb{Z}_{2^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$. It is easy to see that

$$A\mathrm{E}(\lambda^j) = \mathrm{E}(\lambda^{rj}), \qquad \forall\ j \in \mathbb{Z}_{2^n}; \tag{3.4}$$

i.e., A permutes the eigen-subspaces $\mathrm{E}(\lambda^j)$ for $j \in \mathbb{Z}_{2^n}$ the same as μ_r permutes $\mathbb{Z}_{2^n}$, cf. [6, Lemma 2.5]. For any μ_r-orbit $\Gamma_k = \{j_k, rj_k, \cdots, r^{2^b-1}j_k\}$ on $\mathbb{Z}_{2^n}$ of length 2^b ($b \le a$ of course), the subset $\mathcal{E}_k = \{e_{j_k}, e_{rj_k}, \cdots, e_{r^{2^b-1}j_k}\}$ of $\mathcal{E}$ is a basis of the subspace $V_k = \bigoplus_{i=0}^{2^b-1} \mathrm{E}(\lambda^{r^i j_k})$. And there are $\alpha_1, \cdots, \alpha_{2^b} \in \mathbb{C}$ such that $Ae_{j_k} = \alpha_1 e_{rj_k}$, $Ae_{rj_k} = \alpha_2 e_{r^2 j_k}$, $\cdots$, $Ae_{r^{2^b-1}j_k} = \alpha_{2^b} e_{j_k}$. That is, V_k is invariant by A, and the matrix of A with respect to the basis $\mathcal{E}_k$ is

$$A|_{\mathcal{E}_k} = \begin{pmatrix} 0 & \cdots & 0 & \alpha_{2^b} \\ \alpha_1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & \alpha_{2^b-1} & 0 \end{pmatrix}_{2^b \times 2^b}. \tag{3.5}$$

If $\mathrm{ord}(A) = 2^{a'}$ ($a' \ge a$ of course), then $\omega_{2^b} = \alpha_1 \cdots \alpha_{2^b}$ being a $2^{a'-b}$-th root of unity. Replacing $\mathcal{E}_k$ by

$$\mathcal{E}_k = \{e_{j_k}, Ae_{j_k}, \cdots, A^{2^b-1}e_{j_k}\}, \tag{3.6}$$

we get

$$A|_{\mathcal{E}_k} = \begin{pmatrix} 0 & \cdots & 0 & \omega_{2^b} \\ 1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}_{2^b \times 2^b}. \tag{3.7}$$

In particular, $A|_{\mathcal{E}_k}$ is a permutation matrix once $a' = b$ (because $\omega_{2^b} = 1$ once $a' = b$); in that case, A permutes the set $\mathcal{E}_k$ cyclically.

Similarly to $\langle C \rangle$, by $\langle A, C \rangle$ we denote the group generated by A and C.

Lemma 3.1 ([5, Proposition 2.3], [6, Proposition 2.6])**.** *Let notations be as above. If there is a choice of the basis $\mathcal{E}$ in Eqn* (3.3) *such that the matrix $A|_{\mathcal{E}}$ is a permutation matrix, then the matrix group $\langle A, C\rangle$ is a permutation matrix group.*

The statement "the matrix $A|_{\mathcal{E}}$ is a permutation matrix" is equivalent to saying that "the set $\mathcal{E}$ is A-stable", i.e., A permutes the elements of the set $\mathcal{E}$. In that case we call $\mathcal{E}$ an *A-permutation basis*. The rest of this section contributes mainly to finding such a basis $\mathcal{E}$.

We partition $\mathbb{Z}_{2^n}$ into a disjoint union $\mathbb{Z}_{2^n} = 2\mathbb{Z}_{2^n} \bigcup \mathbb{Z}_{2^n}^*$, where

$$2\mathbb{Z}_{2^n} = \{2k \pmod{2^n} \mid k \in \mathbb{Z}_{2^n}\} = \{0, 2, \cdots, 2(2^{n-1}-1)\},$$

$$\mathbb{Z}_{2^n}^* = 1 + 2\mathbb{Z}_{2^n} = \{1 + 2k \pmod{2^n} \mid k \in \mathbb{Z}_{2^n}\} = \{1, 3, \cdots, 2^n - 1\}.$$

Correspondingly, the basis $\mathcal{E}$ in Eqn (3.3) is partitioned into $\mathcal{E} = \mathcal{E}^2 \cup \mathcal{E}^*$ where $\mathcal{E}^2 = \{e_j \,|\, j \in 2\mathbb{Z}_{2^n}\}$ and $\mathcal{E}^* = \{e_j \,|\, j \in \mathbb{Z}_{2^n}^*\}$. And by Eqn (3.2), $\mathbb{C}^{2^n}$ is decomposed into a direct sum of two subspaces: $\mathbb{C}^{2^n} = V^2 \oplus V^*$ where

$$\begin{aligned} V^2 &= \bigoplus_{j \in 2\mathbb{Z}_{2^n}} \mathrm{E}(\lambda^j) = \bigoplus_{e \in \mathcal{E}^2} \mathbb{C}e, \\ V^* &= \bigoplus_{j \in \mathbb{Z}_{2^n}^*} \mathrm{E}(\lambda^j) = \bigoplus_{e \in \mathcal{E}^*} \mathbb{C}e. \end{aligned} \tag{3.8}$$

Both $2\mathbb{Z}_{2^n}$ and $\mathbb{Z}_{2^n}^*$ are μ_r-invariant. So, for any k with $0 \le k < 2^n$, both V^2 and V^* are AC^k-invariant subspaces of $\mathbb{C}^{2^n}$. By $AC^k|_{V^2}$ and $AC^k|_{V^*}$ we denote the linear transformations of AC^k restricted to V^2 and V^* respectively. Correspondingly, $AC^k|_{\mathcal{E}^2}$ and $AC^k|_{\mathcal{E}^*}$ are matrices of $AC^k|_{V^2}$ and $AC^k|_{V^*}$ respectively.

Remark 3.2. The two elements $0, 2^{n-1} \in \mathbb{Z}_{2^n}$ are always fixed by μ_r for any $r \in \mathbb{Z}_{2^n}^*$. So, the two eigen-subspaces $\mathrm{E}(\lambda^0) = \mathrm{E}(1)$ and $\mathrm{E}(\lambda^{2^{n-1}}) = \mathrm{E}(-1)$ are always A-invariant subspaces.

(i). Note that: *if $\langle A, C\rangle$ is a permutation-like matrix group, then $Ae_0 = e_0$ for any $e_0 \in \mathrm{E}(\lambda^0)$.* Because: in that case $\sum_{G \in \langle A,C\rangle} G \neq 0$ (since the traces of permutation matrices are non-negative integers, and the trace of I is $2^n > 0$), there is a vector $f \in \mathbb{C}^{2^n}$ such that $f_0 = \sum_{G \in \langle A,C\rangle} Gf \neq 0$; it is easy to see that $Cf_0 = f_0$ and $Af_0 = f_0$; hence $f_0 \in \mathrm{E}(\lambda^0)$, and $Ae_0 = e_0$ for any $e_0 \in \mathrm{E}(\lambda^0)$.

(ii). It is obvious that, *if $A|_{\mathcal{E}}$ is a permutation matrix, then $Ae_{2^{n-1}} = e_{2^{n-1}}$.*

Lemma 3.3. *Assume that $A, C \in \mathrm{GL}_{2^n}(\mathbb{C})$, C is a maximal cycle and $A^{-1}CA = C^r$ where $r \in \mathbb{Z}_{2^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$. If $A^{2^a} = I$, then there is a choice of the basis $\mathcal{E}^*$ of V^* in Eqn* (3.8) *such that $A|_{\mathcal{E}^*}$ is a permutation matrix.*

Proof. Since $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$, the length of any μ_r-orbit on $\mathbb{Z}_{2^n}^*$ equals to 2^a. Note that $\mathrm{ord}(A) = 2^a$. The conclusion follows from Eqn (3.7). □

The following lemma exhibits the information for the case when $r = -1$, which is enough for our later quotations.

Lemma 3.4. *Assume that $A, C \in \mathrm{GL}_{2^n}(\mathbb{C})$, C is a maximal cycle and $A^{-1}CA = C^{-1}$. If $\mathcal{G} = \langle A, C\rangle$ is a permutation-like matrix group, then $A^2 = I$ (i.e., $\mathcal{G}$ is a dihedral group of order 2^{n+1}) and one of the following two holds:*

(i) *$Ae_{2^{n-1}} = e_{2^{n-1}}$ for $e_{2^{n-1}} \in \mathrm{E}(\lambda^{2^{n-1}})$, and*

$$\mathrm{char}_{AC^k}(x) = \begin{cases} (x-1)^2(x^2-1)^{2^{n-1}-1}, & k \text{ is even;} \\ (x^2-1)^{2^{n-1}}, & k \text{ is odd.} \end{cases}$$

In that case, there is a choice of the basis $\mathcal{E}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix.

(ii) *$Ae_{2^{n-1}} = -e_{2^{n-1}}$ for $e_{2^{n-1}} \in \mathrm{E}(\lambda^{2^{n-1}})$, and*

$$\mathrm{char}_{AC^k}(x) = \begin{cases} (x^2-1)^{2^{n-1}}, & k \text{ is even;} \\ (x-1)^2(x^2-1)^{2^{n-1}-1}, & k \text{ is odd.} \end{cases}$$

Proof. It is easy to see that $\mathbb{Z}_{2^n}$ is partitioned in to μ_{-1}-orbits as follows:

$$2\mathbb{Z}_{2^n} = \{0\} \cup \{2^{n-1}\} \cup \{2, -2\} \cup \cdots \cup \{2^{n-1}-2,\ -2^{n-1}+2\},$$
$$\mathbb{Z}_{2^n}^* = \{1, -1\} \cup \{3, -3\} \cup \cdots \cup \{2^{n-1}-1,\ -2^{n-1}+1\}.$$

Correspondingly, C is diagonally blocked as follows:

$$\begin{aligned} C|_{\mathcal{E}} &= C|_{\mathcal{E}^2} \oplus C|_{\mathcal{E}^*} \\ &= (1 \oplus -1 \oplus \lambda^2 \oplus \lambda^{-2} \oplus \cdots) \oplus (\lambda^1 \oplus \lambda^{-1} \oplus \lambda^3 \oplus \lambda^{-3} \oplus \cdots). \end{aligned}$$

By Lemma 2.5, either $A^2 = I$ or $A^2 = C^{2^{n-1}}$. We prove that $A^2 \neq C^{2^{n-1}}$ by contradiction. Suppose that $A^2 = C^{2^{n-1}}$, i.e., $A|_{\mathcal{E}} = A|_{\mathcal{E}^*} \oplus A|_{\mathcal{E}^2}$ with

$$A^2|_{\mathcal{E}^2} = I_{2^{n-1}}, \qquad A^2|_{\mathcal{E}^*} = -I_{2^{n-1}}.$$

By Eqn (3.7), After a suitable choice of the basis $\mathcal{E}$ we can assume that

$$A|_{\mathcal{E}^2} = 1 \oplus \pm 1 \oplus \overbrace{P \oplus P \oplus \cdots \oplus P}^{2^{n-2}-1} \qquad \text{where } P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

and

$$A|_{\mathcal{E}^*} = \overbrace{Q \oplus Q \oplus \cdots \oplus Q}^{2^{n-2}} \qquad \text{where } Q = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

So, $\mathrm{char}_{A|_{\mathcal{E}^2}}(x) = (x-1)(x \mp 1)(x^2-1)^{2^{n-2}-1}$ and $\mathrm{char}_{A|_{\mathcal{E}^*}}(x) = (x^2+1)^{2^{n-2}}$. We get

$$\mathrm{char}_A(x) = (x-1)(x \mp 1)(x^2+1)(x^4-1)^{2^{n-2}-1}.$$

By Lemma 2.7, the second factor of the right hand side has to be $x+1$. Hence

$$A|_{\mathcal{E}} = A|_{\mathcal{E}^2} \oplus A|_{\mathcal{E}^*} = (1 \oplus -1 \oplus P \oplus P \oplus \cdots) \oplus (Q \oplus Q \oplus \cdots).$$

Then we get the matrix of AC as follows:

$$AC = \left(1 \oplus 1 \oplus \begin{pmatrix} & \lambda^2 \\ \lambda^{-2} & \end{pmatrix} \oplus \cdots\right) \oplus \left(\begin{pmatrix} & -\lambda^{-1} \\ \lambda & \end{pmatrix} \oplus \begin{pmatrix} & -\lambda^{-3} \\ \lambda^3 & \end{pmatrix} \oplus \cdots\right)$$

and the characteristic polynomial as follows:

$$\begin{aligned} \mathrm{char}_{AC}(x) &= (x-1)^2(x^2-1)^{2^{n-2}-1}(x^2+1)^{2^{n-2}} \\ &= (x-1)^2(x^2+1)(x^4-1)^{2^{n-2}-1}. \end{aligned}$$

By Lemma 2.7, AC is not similar to a permutation matrix. However, the assumption of the lemma says that AC is similar to a permutation matrix. The contradiction forces that $A^2 \neq C^{2^{n-1}}$.

Thus, we obtain that $A^2 = I$ and

$$A|_{\mathcal{E}^*} = \overbrace{P \oplus P \oplus \cdots \oplus P}^{2^{n-2}}, \qquad A|_{\mathcal{E}} = 1 \oplus \pm 1 \oplus \overbrace{P \oplus P \oplus \cdots \oplus P}^{2^{n-2}-1}.$$

It is easy to check that:

- if $A|_{\mathcal{E}} = 1 \oplus 1 \oplus \overbrace{P \oplus \cdots \oplus P}^{2^{n-1}-1}$, then (i) holds;
- otherwise, $A|_{\mathcal{E}} = 1 \oplus -1 \oplus \overbrace{P \oplus \cdots \oplus P}^{2^{n-1}-1}$ and (ii) holds. □

In the following we consider the case when $r \not\equiv -1 \pmod{2^n}$ and find a basis $\mathcal{E}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix. We start it with an investigation of the characteristic polynomial of AC^k.

Lemma 3.5. *Let $A, C \in \mathrm{GL}_{2^n}(\mathbb{C})$. Assume that C is a maximal cycle, $A^{-1}CA = C^r$ where $-1 \neq r \in \mathbb{Z}_{2^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$, and $A^{2^a} = I$. Let V^2 and V^* be the subspaces in Eqn (3.8).*

(i) *If $r = 1 + 2^{n-a}v$ where $0 \leq a \leq n-2$ and v is odd, then*

$$\mathrm{char}_{(AC^k)|_{V^*}}(x) = \begin{cases} \Phi_{2^{n-\nu_2(k)}}(x)^{2^{\nu_2(k)}}, & 0 \leq \nu_2(k) < n-a; \\ (x^{2^a}-1)^{2^{n-a-1}}, & \nu_2(k) \geq n-a, \text{ or } k = 0. \end{cases}$$

(ii) *If $r = -1 + 2^{n-a}v$ where $1 \leq a \leq n-2$ and v is odd, then*

$$\mathrm{char}_{(AC^k)|_{V^*}}(x) = \begin{cases} (x^{2^a}+1)^{2^{n-a-1}} = \Phi_{2^{a+1}}(x)^{2^{n-a-1}}, & k \text{ is odd}; \\ (x^{2^a}-1)^{2^{n-a-1}}, & k \text{ is even}. \end{cases}$$

Proof. By the assumption, $\mathbb{Z}_{2^n}^*$ is partitioned into μ_r-orbits $\Gamma_1, \cdots, \Gamma_h$ with the lengths all being 2^a and the number $h = 2^{n-a-1}$:

$$\Gamma_1 = \{j_1, rj_1, \cdots, r^{2^a-1}j_1\}, \ \cdots, \ \Gamma_h = \{j_h, rj_h, \cdots, r^{2^a-1}j_h\}.$$

By Eqn (3.3) and Eqn (3.6), we can assume that $\mathcal{E}^* = \mathcal{E}_1^* \cup \cdots \cup \mathcal{E}_h^*$ with

$$\mathcal{E}_i^* = \left\{e_{j_i}, \ Ae_{j_i}, \ \cdots, \ A^{r^{2^a-1}}e_{j_i}\right\}, \qquad i = 1, \cdots, h. \tag{3.9}$$

Denote the matrices of A and C restricted to $\mathcal{E}_i^*$ by $A_i = A|_{\mathcal{E}_i^*}$ and $C_i = C|_{\mathcal{E}_i^*}$ for $i = 1, \cdots, h$. Since $A_i^{2^a} = I$, by Lemma 3.3 we have

$$A_i = \begin{pmatrix} 0 & \cdots & 0 & 1 \\ 1 & \ddots & \ddots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 0 \end{pmatrix}_{2^a \times 2^a}, \ C_i^k = \begin{pmatrix} \lambda^{kj_i} & & & \\ & \lambda^{kj_i r} & & \\ & & \ddots & \\ & & & \lambda^{kj_i r^{2^a-1}} \end{pmatrix}_{2^a \times 2^a}.$$

Thus $AC^k|_{\mathcal{E}^*} = \bigoplus_{i=1}^h A_i C_i^k$, and $\mathrm{char}_{A_i C_i^k}(x) = x^{2^a} - \lambda^{j_i k(1+r+\cdots+r^{2^a-1})}$. The conclusion is obviously true if $k = 0$. So we further assume that $k \neq 0$.

Writing $k = k'2^{\nu_2(k)}$ with k' being odd, and writing $1+r+\cdots+r^{2^a-1} = a'2^{a''}$ with a' being odd and $a'' = \nu_2(1 + r + \cdots + r^{2^a-1})$, we have

$$\mathrm{char}_{A_i C_i^k}(x) = x^{2^a} - \lambda^{j_i a' k' 2^{a''+\nu_2(k)}}, \qquad i = 1, \cdots, h.$$

If $a'' + \nu_2(k) \geq n$, then $\lambda^{j_i a' k' 2^{a''+\nu_2(k)}} = 1$. However, if $a'' + \nu_2(k) < n$, then $\lambda^{j_i a' k' 2^{a''+\nu_2(k)}}$ is a primitive $2^{n-a''-\nu_2(k)}$-th root of unity, and, by Lemma 2.1 (ii), the collection of

$$\lambda^{j_i a' k' 2^{a''+\nu_2(k)}}, \qquad i = 1, \cdots, h,$$

is just the collection of all primitive $2^{n-a''-\nu_2(k)}$-th roots of unity, each of which appears with multiplicity $\frac{h}{2^{n-a''-\nu_2(k)-1}} = 2^{a''-a+\nu_2(k)}$. By Eqn (2.1), we obtain that

$$\mathrm{char}_{(AC^k)|_{V^*}}(x) = \begin{cases} (x^{2^a}-1)^{n-a-1}, & a'' + \nu_2(k) \geq n; \\ \Phi_{2^{n+a-a''-\nu_2(k)}}(x)^{2^{a''-a+\nu_2(k)}}, & a'' + \nu_2(k) < n. \end{cases}$$

If $r = 1 + 2^{n-a}v$ where $0 \leq a \leq n-2$ and v is odd, then, by Lemma 2.4(ii), $a'' = a$; hence (i) is proved.

Next, assume that $r = -1 + 2^{n-a}v$ where $1 \leq a \leq n-2$ and v is odd. By Lemma 2.4(iii), $a'' = n-1$. So $a'' + \nu_2(k) < n$ if and only if k is odd; in that case, $n + a - a'' - \nu_2(k) = a + 1$. Noting that $\Phi_{2^{a+1}}(x) = x^{2^a} + 1$, we are done for the conclusion (ii). □

The second step is a preparation for induction. Let $r \in \mathbb{Z}_{2^n}^*$. Then μ_r (the action by multiplication) acts on both $\mathbb{Z}_{2^{n-1}}$ and $2\mathbb{Z}_{2^n}$, and the map

$$\mathbb{Z}_{2^{n-1}} \longrightarrow 2\mathbb{Z}_{2^n}, \ \ t \longmapsto 2t, \tag{3.10}$$

is an isomorphism between the μ_r-acted sets $\mathbb{Z}_{2^{n-1}}$ and $2\mathbb{Z}_{2^n}$.

Lemma 3.6. *Let notation be as in Lemma 3.5. Assume further that the matrix group $\langle A, C\rangle$ is permutation-like. Then $\langle A|_{V^2}, C|_{V^2}\rangle$ is a permutation-like matrix group of dimension 2^{n-1} with $C|_{V^2}$ being a maximal cycle.*

Proof. Since $C|_{\mathcal{E}^2} = \bigoplus_{j=0}^{2^{n-1}-1} \lambda^j$, with the isomorphism (3.10), $C|_{V^2}$ is a maximal cycle of length 2^{n-1}. Note that $\langle A, C\rangle = \{A^\ell C^k \mid 0 \le \ell < 2^a,\ 0 \le k < 2^n\}$. To complete the proof, it is enough to show that any $(A^\ell C^k)|_{V^2}$ is similar to a permutation matrix.

Let $A' = A^\ell$, $a' = a - \nu_2(\ell)$ and $r' = r^\ell$. Then $A'^{-1}CA' = C^{r'}$ and $\mathrm{ord}(A') = \mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r') = 2^{a'}$. We compute $\mathrm{char}_{(A^\ell C^k)|_{V^2}}(x) = \mathrm{char}_{(A'C^k)|_{V^2}}(x)$ in two cases.

Case 1: $r \in \langle 5\rangle$ or $\nu_2(\ell) \ge 1$ (cf. Remark 2.2). We can use Lemma 3.5(i) to $A'C^k$ and get that

$$\mathrm{char}_{(A^\ell C^k)|_{V^*}}(x) = \begin{cases} \Phi_{2^{n-\nu_2(k)}}(x)^{2^{\nu_2(k)}}, & 0 \le \nu_2(k) < n - a'; \\ (x^{2^{a'}} - 1)^{2^{n-a'-1}}, & \nu_2(k) \ge n - a',\ \text{or } k = 0. \end{cases}$$

Since $A^\ell C^k$ is similar to a permutation matrix, by Lemma 2.7 and its corollary,

$$\mathrm{char}_{(A^\ell C^k)|_{V^2}}(x) = \begin{cases} (x^{2^{n-\nu_2(k)-1}} - 1)^{2^{\nu_2(k)}}, & 0 \le \nu_2(k) < n - a'; \\ \prod_i (x^{2^i} - 1)^{j_i}, & \nu_2(k) \ge n - a',\ \text{or } k = 0. \end{cases} \tag{3.11}$$

Case 2: $r = -1 + 2^{n-a}v$ with v being odd and $\nu_2(\ell) = 0$ (i.e. $a' = a$). By Lemma 3.5(ii) (applied to $A'C^k$),

$$\mathrm{char}_{(A^\ell C^k)|_{V^*}}(x) = \begin{cases} \Phi_{2^{a+1}}(x)^{2^{n-a-1}}, & k \text{ is odd}; \\ (x^{2^a} - 1)^{2^{n-a-1}}, & k \text{ is even}. \end{cases}$$

Since $A^\ell C^k$ is similar to a permutation matrix, by Lemma 2.7 and its corollary,

$$\mathrm{char}_{A^\ell C^k|_{V^2}}(x) = \begin{cases} (x^{2^a} - 1)^{2^{n-a-1}}, & k \text{ is odd}; \\ \prod_i (x^{2^i} - 1)^{j_i}, & k \text{ is even}. \end{cases} \tag{3.12}$$

In both cases, by Lemma 2.7 again, we can conclude that $A^\ell C^k|_{V^2}$ is similar to a permutation matrix. □

Now we can get enough information for $A|_{\mathcal{E}}$ when $r \not\equiv -1 \pmod{2^n}$. We proceed in two cases: $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2$, or $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) > 2$.

Lemma 3.7. *Let $A, C \in \mathrm{GL}_{2^n}(\mathbb{C})$. Assume that C is a maximal cycle, $A^{-1}CA = C^r$ where $-1 \ne r \in \mathbb{Z}_{2^n}^*$ with $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2$, and $A^2 = I$. Assume further that $\langle A, C\rangle$ is a permutation-like matrix group.*

(i) *If* $r = 1 + 2^{n-1}$, *then* $A|_{V^2} = I$.

(ii) *If* $r = -1 + 2^{n-1}$, *then* $\mathrm{char}_{A|_{V^2}}(x) = (x-1)^2(x^2-1)^{2^{n-1}-1}$.

In the both cases there is a choice of the basis $\mathcal{E}$ *in Eqn* (3.3) *such that* $A|_{\mathcal{E}}$ *is a permutation matrix.*

Proof. By Lemma 3.6, $\langle A|_{V^2}, C|_{V^2}\rangle$ is a permutation-like matrix group of dimension 2^{n-1} with $C|_{V^2}$ being a maximal cycle.

(i). Obviously, $r = 1 + 2^{n-1}$ acts by multiplication on $2\mathbb{Z}_{2^n}$ trivially. So, $A|_{\mathcal{E}^2}$ is a diagonal matrix, hence $A|_{\mathcal{E}^2}$ commutes with $C|_{\mathcal{E}^2}$ which is a maximal cycle of dimension 2^{n-1}. By Lemma 2.6, $A|_{\mathcal{E}^2} \in \langle C|_{\mathcal{E}^2}\rangle$. But $(A|_{\mathcal{E}^2})^2 = I$ and $\mathrm{ord}(C|_{\mathcal{E}^2}) = 2^{n-1}$. Thus $A|_{\mathcal{E}^2} = I_{2^{n-1}\times 2^{n-1}}$ or $= (C|_{\mathcal{E}^2})^{2^{n-2}}$. Suppose that $A|_{\mathcal{E}^2} = (C|_{\mathcal{E}^2})^{2^{n-2}}$, i.e., $(A|_{\mathcal{E}^2})(C|_{\mathcal{E}^2})^{-2^{n-2}} = I_{2^{n-1}\times 2^{n-1}}$. Then

$$\mathrm{char}_{(A|_{\mathcal{E}^2})(C|_{\mathcal{E}^2})^{-2^{n-2}}}(x) = (x-1)^{2^{n-1}}.$$

However, $\nu_2(-2^{n-2}) = n-2$, and by Eqn (3.11),

$$\mathrm{char}_{(A|_{\mathcal{E}^2})(C|_{\mathcal{E}^2})^{-2^{n-2}}}(x) = (x^2-1)^{2^{n-2}}.$$

We reached an impossible equality "$(x-1)^{2^{n-1}} = (x^2-1)^{2^{n-2}}$". So it is impossible that $A|_{\mathcal{E}^2} = (C|_{\mathcal{E}^2})^{2^{n-2}}$, and it has to be the case that $A|_{\mathcal{E}^2} = I_{2^{n-1}\times 2^{n-1}}$. By Lemma 3.3, we already have a basis $\mathcal{E}^*$ of V^* such that $A|_{\mathcal{E}^*}$ is a permutation matrix. Thus the basis $\mathcal{E}$ in Eqn (3.3) can be obtained such that $A|_{\mathcal{E}}$ is a permutation matrix.

(ii). Note that $r = -1 + 2^{n-1} \equiv -1 \pmod{2^{n-1}}$ (cf. Eqn (3.10)), so

$$(A|_{V^2})^{-1}(C|_{V^2})(A|_{V^2}) = (C|_{V^2})^{-1}.$$

From Lemma 3.6(ii) and its Eqn (3.12), we have seen that $\langle A|_{\mathcal{E}^2}, C|_{\mathcal{E}^2}\rangle$ is a permutation-like matrix group of dimension 2^{n-1} with $C|_{\mathcal{E}^2}$ being a maximal cycle, and

$$\mathrm{char}_{AC^k|_{V^2}}(x) = \begin{cases} (x^2-1)^{2^{n-2}}, & k \text{ is odd;} \\ \prod_i (x^{2^i}-1)^{j_i}, & k \text{ is even.} \end{cases}$$

Comparing it with Lemma 3.4, we see that $(A|_{V^2})e_{2^{n-1}} = e_{2^{n-1}}$, and, with a suitable choice of the basis $\mathcal{E}^2$, $A|_{\mathcal{E}^2}$ is a permutation matrix. □

Lemma 3.8. *Assume thar* $A, C \in \mathrm{GL}_{2^n}(\mathbb{C})$, C *is a maximal cycle and* $A^{-1}CA = C^r$ *where* $r \in \mathbb{Z}_{2^n}^*$ *with* $\mathrm{ord}_{\mathbb{Z}_{2^n}^*}(r) = 2^a$. *If* $a \geq 2$ *and* $A^{2^a} = I$, *then there is a choice of the basis* $\mathcal{E}$ *in Eqn* (3.3) *such that* $A|_{\mathcal{E}}$ *is a permutation matrix.*

Proof. We prove it by induction on n. By Remark 2.2,

$$r = \pm 1 + 2^{n-a}v, \quad \text{with } v \text{ being odd and } 2 \leq a \leq n-2.$$

From Lemma 3.3, we have a basis $\mathcal{E}^*$ of V^* in Eqn (3.8) such that $A|_{\mathcal{E}^*}$ is a permutation matrix. By Lemma 3.6, $\langle A|_{V^2}, C|_{V^2}\rangle$ is a permutation-like matrix group of dimension 2^{n-1} with $C|_{V^2}$ being a maximal cycle. Let $A' = A^{2^{a-1}}$. Then $A'^2 = I$ and $A'^{-1}CA' = C^{r^{2^{a-1}}}$. Since 2^{a-1} is even, we see that $r^{2^{a-1}} \in \langle 5\rangle$, hence $r^{2^{a-1}}$ is the unique element of order 2 in $\langle 5\rangle$, i.e., $r^{2^{a-1}} \equiv 1+2^{n-1} \pmod{2^n}$; see Remark 2.2. By Lemma 3.7(i), $A'|_{V^2} = I$; i.e.,

$$(A|_{V^2})^{2^{a-1}} = A^{2^{a-1}}|_{V^2} = I.$$

Note that $\mathrm{ord}_{\mathbb{Z}^*_{2^{n-1}}}(r) = 2^{a-1}$. By induction, we have a basis $\mathcal{E}^2$ of V^2 in Eqn (3.8) such that $A|_{\mathcal{E}^2}$ is a permutation matrix. Then $\mathcal{E} = \mathcal{E}^* \cup \mathcal{E}^2$ is a basis of $\mathbb{C}^{2^n}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix. □

Theorem 3.9. *Assume that $\mathcal{G}$ is a 2^n-dimensional permutation-like matrix group which contains a maximal cycle C such that $\langle C\rangle$ is normal in $\mathcal{G}$ and the quotient group $\mathcal{G}/\langle C\rangle$ is cyclic. Then $\mathcal{G}$ is a permutation matrix group.*

Proof. By the assumption we can assume that $\mathcal{G} = \langle A, C\rangle$ and $A^{-1}CA = C^r$ for an $r \in \mathbb{Z}^*_{2^n}$ with $\mathrm{ord}_{\mathbb{Z}^*_{2^n}}(r) = 2^a$ where $0 \le a \le n-2$. By Lemma 2.6, $\langle C\rangle$ is self-centralized in $\mathcal{G}$. By Lemma 2.5 and Lemma 3.4, we can further assume that $A^{2^a} = I$ and $1 \le a \le n-2$. By Remark 2.2, we prove the theorem in three cases.

Case 1: $r = -1$, hence $a = 1$ and $A^2 = I$. By Lemma 3.4, $Ae_{2^{n-1}} = \pm e_{2^{n-1}}$. If $Ae_{2^{n-1}} = -e_{2^{n-1}}$, then we replace A by AC. So we can assume that $Ae_{2^{n-1}} = e_{2^{n-1}}$. By Lemma 3.4 (i), there is a basis $\mathcal{E}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix. By Lemma 3.1, $\mathcal{G}$ is a permutation matrix group.

Case 2: $r = \pm 1 + 2^{n-1}$ (hence $a = 1$). By Lemma 2.5(ii), we can assume that $A^2 = I$. By Lemma 3.7, there is a basis $\mathcal{E}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix. By Lemma 3.1, $\mathcal{G}$ is a permutation matrix group.

Case 3: $r = \pm 1 + 2^{n-a}v$ with v being odd and $2 \le a \le n-2$. By Lemma 2.5(ii), we can assume that $A^{2^a} = I$. By Lemma 3.8, there is a basis $\mathcal{E}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix. By Lemma 3.1, we are done. □

4 The quotient $\mathcal{G}/\langle C\rangle$ is non-cyclic

In this section we consider any 2^n-dimensional permutation-like matrix group $\mathcal{G}$ with a maximal cycle C such that $\langle C\rangle$ is normal in $\mathcal{G}$ and the quotient group $\mathcal{G}/\langle C\rangle$ is not cyclic, and prove that $\mathcal{G}$ is in fact a permutation matrix group.

It is easy to extend Lemma 3.1 as follows.

Lemma 4.1. *Let $A, B, C \in \mathrm{GL}_{2^n}(\mathbb{C})$ with C being a maximal cycle and the subgroup $\langle C\rangle$ being normal in $\mathcal{G} = \langle A, B, C\rangle$. If there is a choice of the basis $\mathcal{E}$ in Eqn (3.3) such that both $A|_{\mathcal{E}}$ and $B|_{\mathcal{E}}$ are permutation matrices, then the matrix group $\mathcal{G}$ is a permutation matrix group.*

Proof. Let $f = \sum_{e\in\mathcal{E}} e$ and $\mathcal{F} = \{f,\ Cf,\ , C^2f,\ \cdots,\ C^{2^n-1}f\}$. Because the matrix to transform the basis $\mathcal{E}$ to the set $\mathcal{F}$ is a Vandemond matrix formed by $1, \lambda, \cdots, \lambda^{2^n-1}$ which is non-degenerate, $\mathcal{F}$ is a basis of $\mathbb{C}^{2^n}$. With respect to the basis $\mathcal{F}$, $C|_{\mathcal{F}}$ is obviously a permutation matrix. Since A permutes the elements of $\mathcal{E}$, we have $Af = f$. For any $C^jf \in \mathcal{F}$,

$$A(C^jf) = (AC^jA^{-1})(Af) = C^{r^{-1}j}f \in \mathcal{F}.$$

Thus $A|_{\mathcal{F}}$ is a permutation matrix. Similarly, $B|_{\mathcal{F}}$ is a permutation matrix. So, with respect to the basis $\mathcal{F}$ every element of $\mathcal{G}$ is a permutation matrix. □

Lemma 4.2. *Let $r = 1 + 2^{n-a} \in \mathbb{Z}_{2^n}^*$ where $1 \le a \le n-2$. Then there are $j_1, \cdots, j_m \in \mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$ such that all the μ_r-orbits on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$ are written as follows (note that $b_t = 1$, $1 \le t \le m$, is allowed):*

$$\begin{aligned}
&\Gamma_1 = \{j_1, j_1r, \cdots, j_1r^{2^{b_1}-1}\}, && \mu_{-1}\Gamma_1 = \{-j_1, -j_1r, \cdots, -j_1r^{2^{b_1}-1}\},\\
&\Gamma_2 = \{j_2, j_2r, \cdots, j_2r^{2^{b_2}-1}\}, && \mu_{-1}\Gamma_2 = \{-j_2, -j_2r, \cdots, -j_2r^{2^{b_2}-1}\},\\
&\cdots\quad\cdots && \cdots\quad\cdots\\
&\Gamma_m = \{j_m, j_mr, \cdots, j_mr^{2^{b_m}-1}\}, && \mu_{-1}\Gamma_m = \{-j_m, -j_mr, \cdots, -j_mr^{2^{b_m}-1}\}.
\end{aligned}$$

Proof. For $k \in \mathbb{Z}_{2^n}$, $rk \equiv -k \pmod{2^n}$ if and only if $(r+1)k \equiv 0 \pmod{2^n}$. However, $\nu_2(r+1) = 1$ since $r + 1 = 2 + 2^{n-a}$ and $n - a \ge 2$. We see that $rk \equiv -k \pmod{2^n}$ if and only if $k \equiv 0$ or $2^{n-1} \pmod{2^n}$. So, for any μ_r-orbit $\Gamma = \{j, jr, \cdots, jr^{2^b-1}\}$ on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$, the set $\mu_{-1}\Gamma = \{-j, -jr, \cdots, -jr^{2^b-1}\}$ is another μ_r-orbit on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$, and

$$\mu_{-1}(\mu_{-1}\Gamma) = \mu_{-1}(\{-j, -jr, \cdots, -jr^{2^b-1}\}) = \Gamma.$$

Starting from any μ_r-orbit $\Gamma_1 = \{j_1, j_1r, \cdots, j_1r^{2^{b_1}-1}\}$ on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$, we get a pair $\Gamma_1, \mu_{-1}\Gamma_1$ of μ_r-orbits on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$. Taking any μ_r-orbit $\Gamma_2 = \{j_2, j_2r, \cdots, j_2r^{2^{b_2}-1}\}$ on $\mathbb{Z}_{2^n}\backslash\big(\{0, 2^{n-1}\} \cup \Gamma \cup \mu_{-1}\Gamma\big)$, we get another pair $\Gamma_2, \mu_{-1}\Gamma_2$ on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$. Go on in this way till to $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$ exhausted, we obtain all the μ_r-orbits on $\mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$ as listed pairwise in the lemma. □

Theorem 4.3. *Assume that $\mathcal{G}$ is a 2^n-dimensional permutation-like matrix group which contains a maximal cycle C such that $\langle C\rangle$ is normal and the quotient group $\mathcal{G}/\langle C\rangle$ is not cyclic. Then $\mathcal{G}$ is a permutation matrix group.*

Proof. Since $\langle C\rangle$ is self-centralized in $\mathcal{G}$ (see Lemma 2.6), $\mathcal{G}/\langle C\rangle$ is isomorphic to a subgroup $\mathcal{H}$ of the multiplicative group $\mathbb{Z}_{2^n}^*$. By Lemma 2.3 (ii), $\mathcal{H} = \langle r\rangle \times \langle -1\rangle$ where $r = 1 + 2^{n-a}$ with $1 \le a \le n-2$. Then there are $A, B \in \mathcal{G}$ such that

$$\mathcal{G} = \langle A, B, C\rangle, \quad B^{-1}CB = C^{-1}, \quad A^{-1}CA = C^r.$$

By Lemma 2.5(i) and Lemma 3.4, we further assume that $A^{2^a} = I$ and $B^2 = I$.

First we show that *there is a suitable choice of B such that $AB = BA$.* Since the quotient group $\mathcal{G}/\langle C\rangle \cong \mathcal{H}$ is commutative, in $\mathcal{G}/\langle C\rangle$ the image of $B^{-1}AB$ coincides with the image of A itself. So there is a $C^k \in \langle C\rangle$ such that $B^{-1}AB = AC^k$. Note that $\mathrm{ord}(B^{-1}AB) = \mathrm{ord}(A) = 2^a$, by Eqn (2.4),

$$I = (B^{-1}AB)^{2^a} = (AC^k)^{2^a} = A^{2^a}C^{k(r^{2^a-1}+\cdots+r+1)} = C^{k(r^{2^a-1}+\cdots+r+1)}.$$

By Lemma 2.4(ii), $\nu_2(r^{2^a-1}+\cdots+r+1) = a$. So the above equality implies that $\nu_2(k) \geq n-a$. We can write $k = 2^{n-a}h$. Noting that $r-1 = 2^{n-a}$, we obtain:

$$\begin{aligned}(BC^h)^{-1}A(BC^h) = C^{-h}B^{-1}ABC^h = C^{-h}AC^kC^h = AC^{-rh}C^kC^h\\ = AC^{k+h-rh} = AC^{2^{n-a}h-2^{n-a}h} = A.\end{aligned}$$

Replacing B with BC^h, we still have $\mathcal{G} = \langle A, B, C\rangle$, $B^{-1}CB = C^{-1}$ and $B^2 = I$ (since $(BC^h)^2 = I$, see Lemma 2.5(ii.1)), and we have a further condition that $AB = BA$.

By Lemma 3.7(i) (for $a = 1$) and Lemma 3.8 (for $a > 1$), there exists a basis $\mathcal{E}$ in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix. Specifically, $Ae_0 = e_0$ for any $e_0 \in \mathrm{E}(\lambda^0)$ and $Ae_{2^{n-1}} = e_{2^{n-1}}$ for any $e_{2^{n-1}} \in \mathrm{E}(\lambda^{2^{n-1}})$, see Remark 3.2. Using the notation in Lemma 4.2 and the method in Eqn (3.7), we construct the basis $\mathcal{E}$ precisely by choosing e_j for $j \in \mathbb{Z}_{2^n}\backslash\{0, 2^{n-1}\}$ as follows.

Let Γ_t and $\mu_{-1}\Gamma_t$, for $t = 1, \cdots, m$, be the μ_r-orbits in Lemma 4.2. We take a non-zero vector $e_{j_t} \in \mathrm{E}(\lambda^{j_t})$. By Eqn (3.7), $\mathcal{E}_t = \{e_{j_t}, Ae_{j_t}, \cdots, A^{2^{b_t}-1}e_{j_t}\}$ is a basis of the subspace $V_t = \bigoplus_{i=0}^{2^{b_t}-1}\mathrm{E}(\lambda^{r^i j_t})$ such that $A|_{\mathcal{E}_t}$ is a permutation matrix. Since $Be_{j_t} \in \mathrm{E}(\lambda^{-j_t})$ (see Eqn (3.4)), we can take $e_{-j_t} = Be_{j_t}$. Then $\mathcal{E}'_t = \{e_{-j_t}, Ae_{-j_t}, \cdots, A^{2^{b_t}-1}e_{-j_t}\}$ is a basis of the subspace $V'_t = \bigoplus_{i=0}^{2^{b_t}-1}\mathrm{E}(\lambda^{-r^i j_t})$ such that $A|_{\mathcal{E}'_t}$ is a permutation matrix. Let $\mathcal{E} = \{e_0, e_{2^{n-1}}\}\bigcup_{t=1}^{m}(\mathcal{E}_t \cup \mathcal{E}'_t)$. Then $\mathcal{E}$ is a basis in Eqn (3.3) such that $A|_{\mathcal{E}}$ is a permutation matrix.

On the other hand, $Be_{-j_t} = B^2e_{j_t} = e_{j_t}$. i.e., B interchanges e_{j_t} and e_{-j_t}. Since B commutes with A, we get that $B(Ae_{j_t}) = A(Be_{j_t}) = Ae_{-j_t}$. It is the same that $BAe_{-j_t} = Ae_{j_t}$. So, B interchanges Ae_{j_t} and Ae_{-j_t}. Similarly, B interchanges $A^ie_{j_t}$ and $A^ie_{-j_t}$ for $i = 0, 1, \cdots, 2^{b_{j_t}}-1$. In a word, $B|_{\mathcal{E}_t\cup\mathcal{E}'_t}$ is a permutation matrix. Of course, $Be_0 = e_0$, see Remark 3.2. To complete the proof, it is enough to show that $Be_{2^{n-1}} = e_{2^{n-1}}$; because: it implies that $B|_{\mathcal{E}}$ is a permutation matrix, and by Lemma 4.1, $\mathcal{G}$ is a permutation matrix group.

From an easy computation:

$$(AB)^{-1}C(AB) = B^{-1}(A^{-1}CA)B = B^{-1}C^{1+2^{n-a}}B = C^{-1-2^{n-a}},$$

we have $(AB)^{-1}C(AB) = C^{-1+2^{n-a}v}$ with $v = -1$ being odd. Obviously, $\langle AB, C\rangle$ is a permutation-like matrix group and $(AB)^{2^a} = A^{2^a}B^{2^a} = I$ (since $AB = BA$). By Lemma 3.7(ii) and Lemma 3.8, there is a basis $\mathcal{E}'$ in Eqn (3.3) such that $(AB)|_{\mathcal{E}'}$ is a permutation matrix. In particular, $(AB)e_{2^{n-1}} = e_{2^{n-1}}$, see Remark 3.2. Note that $Ae_{2^{n-1}} = e_{2^{n-1}}$ implies that $A^{-1}e_{2^{n-1}} = e_{2^{n-1}}$. Then $Be_{2^{n-1}} = A^{-1}(ABe_{2^{n-1}}) = A^{-1}e_{2^{n-1}} = e_{2^{n-1}}$. We are done. □

Acknowledgments

The authors are supported by NSFC through the grant number 11271005.

References

[1] J. L. Alperin, R. B. Bell, Groups and Representations, GTM 162, Springer-Verlag, New York, 1997.

[2] Bocong Chen, Hai Q. Dinh, Yun Fan, and San Ling, Polyadic constacyclic codes, IEEE Trans. Inform. Theory 61(2015), 4895-4904.

[3] G. Cigler, Groups of matrices with prescribed spectrum, Doctoral dissertation, 2005, http://matknjiz.si/doktotati/2005/10921-83.pdf

[4] G. Cigler, Permutation-like matrix groups, Linear Algebra and its Applications 422(2007), 486-505.

[5] Guodong Deng, Yun Fan, Permutation-like matrix group with a maximal cycle of prime square length, Linear Algebra and its Applications 450(2014), 44-55.

[6] Guodong Deng, Yun Fan, Permutation-like matrix groups with a maximal cycle of power of odd prime length, Linear Algebra and its Applications 480(2015), 1-10.

[7] D.J.S. Robinson, A Course in Groups Theory, GTM 80, Springer-Verlag, New York, 1980.